

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
6 May 2005 (06.05.2005)

PCT

(10) International Publication Number  
**WO 2005/040114 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 209/30**,  
A61K 31/404, A61P 43/00, C07D 403/12, 401/12, 417/12

Oxagen Limited, 91 Milton Park, Abingdon, Oxon OX14 4RY (GB).

(21) International Application Number:  
PCT/GB2004/004336

(74) Agents: **ROBERTS, Alison, Christine** et al.; Kilburn & Strode, 20 Red Lion Street, London WC1R 4PJ (GB).

(22) International Filing Date: 13 October 2004 (13.10.2004)

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0324084.3 14 October 2003 (14.10.2003) GB  
0400716.7 14 January 2004 (14.01.2004) GB

(71) Applicant (*for all designated States except US*): **OXAGEN LIMITED** [GB/GB]; 91 Milton Park, Abingdon, Oxon OX14 4RY (GB).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MIDDLEMISS, David** [GB/GB]; 62 Thorley Hill, Bishop's Stortford, Herts CM23 3NF (GB). **ASHTON, Mark, Richard** [GB/GB]; Evotec OAI, 151 Milton Park, Abingdon, Oxon OX14 4RY (GB). **BOYD, Edward, Andrew** [GB/GB]; Evotec OAI Limited, 151 Milton Park, Abingdon, OXON OX14 4RY (GB). **BROOKFIELD, Frederick, Arthur** [GB/GB]; Evotec OAI Limited, 151 Milton Park, Abingdon, Oxon OX14 4RY (GB). **ARMER, Richard, Edward** [GB/GB];


(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOUNDS HAVING CRTH2 ANTAGONIST ACTIVITY



(I)

(57) Abstract: Compounds of general formula (I) and their pharmaceutically acceptable salts, hydrates, solvates, complexes and prodrugs are useful in the treatment of allergic diseases such as asthma, allergic rhinitis and atopic dermatitis.



WO 2005/040114 A1

### COMPOUNDS HAVING CRTH2 ANTAGONIST ACTIVITY

The present invention relates to compounds which are useful as pharmaceuticals, to methods for preparing these compounds, compositions containing them and their use  
5 in the treatment and prevention of allergic diseases such as asthma, allergic rhinitis and atopic dermatitis and other inflammatory diseases mediated by prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) acting at the CRTH2 receptor on cells including eosinophils, basophils and Th2 lymphocytes.

10 PGD<sub>2</sub> is an eicosanoid, a class of chemical mediator synthesised by cells in response to local tissue damage, normal stimuli or hormonal stimuli or *via* cellular activation pathways. Eicosanoids bind to specific cell surface receptors on a wide variety of tissues throughout the body and mediate various effects in these tissues. PGD<sub>2</sub> is known to be produced by mast cells, macrophages and Th2 lymphocytes and has  
15 been detected in high concentrations in the airways of asthmatic patients challenged with antigen (Murray *et al*, (1986), *N. Engl. J. Med.* **315**: 800-804). Instillation of PGD<sub>2</sub> into airways can provoke many features of the asthmatic response including bronchoconstriction (Hardy *et al*, (1984) *N. Engl. J. Med.* **311**: 209-213; Sampson *et al*, (1997) *Thorax* **52**: 513-518) and eosinophil accumulation (Emery *et al*, (1989) *J. Appl. Physiol.* **67**: 959-962).  
20

The potential of exogenously applied PGD<sub>2</sub> to induce inflammatory responses has been confirmed by the use of transgenic mice overexpressing human PGD<sub>2</sub> synthase which exhibit exaggerated eosinophilic lung inflammation and Th2 cytokine  
25 production in response to antigen (Fujitani *et al*, (2002) *J. Immunol.* **168**: 443-449).

The first receptor specific for PGD<sub>2</sub> to be discovered was the DP receptor which is linked to elevation of the intracellular levels of cAMP. However, PGD<sub>2</sub> is thought to mediate much of its proinflammatory activity through interaction with a G protein-  
30 coupled receptor termed CRTH2 (chemoattractant receptor-homologous molecule expressed on Th2 cells) which is expressed by Th2 lymphocytes, eosinophils and


basophils (Hirai *et al*, (2001) *J. Exp. Med.* **193**: 255-261, and EP0851030 and EP-A-1211513 and Bauer *et al*, EP-A-1170594). It seems clear that the effect of PGD<sub>2</sub> on the activation of Th2 lymphocytes and eosinophils is mediated through CRTH2 since the selective CRTH2 agonists 13,14 dihydro-15-keto-PGD<sub>2</sub> (DK-PGD<sub>2</sub>) and 15R-methyl-PGD<sub>2</sub> can elicit this response and the effects of PGD<sub>2</sub> are blocked by an anti-CRTH2 antibody (Hirai *et al*, 2001; Monneret *et al*, (2003) *J. Pharmacol. Exp. Ther.* **304**: 349-355). In contrast, the selective DP agonist BW245C does not promote migration of Th2 lymphocytes or eosinophils (Hirai *et al*, 2001; Gervais *et al*, (2001) *J. Allergy Clin. Immunol.* **108**: 982-988). Based on this evidence, antagonising PGD<sub>2</sub> at the CRTH2 receptor is an attractive approach to treat the inflammatory component of Th2-dependent allergic diseases such as asthma, allergic rhinitis and atopic dermatitis.

EP-A-1170594 suggests that the method to which it relates can be used to identify compounds which are of use in the treatment of allergic asthma, atopic dermatitis, allergic rhinitis, autoimmune disease, reperfusion injury and a number of inflammatory conditions, all of which are mediated by the action of PGD<sub>2</sub> at the CRTH2 receptor.

Compounds which bind to CRTH2 are taught in WO-A-03066046 and WO-A-03066047. These compounds are not new but were first disclosed, along with similar compounds, in GB 1356834, GB 1407658 and GB 1460348, where they were said to have anti-inflammatory, analgesic and antipyretic activity. WO-A-03066046 and WO-A-03066047 teach that the compounds to which they relate are modulators of CRTH2 receptor activity and are therefore of use in the treatment or prevention of obstructive airway diseases such as asthma, chronic obstructive pulmonary disease (COPD) and a number of other diseases including various conditions of bones and joints, skin and eyes, GI tract, central and peripheral nervous system and other tissues as well as allograft rejection. The compounds described in these documents are indoles with a carboxylic acid group is at the 3-position of the indole ring system a quinoline, quinazoline or benzothiazole group at the 1-position.

The present invention relates to novel compounds which bind to CRTH2 and which will therefore also be useful in the treatment of diseases and conditions mediated by the activity of PGD<sub>2</sub> at the CRTH2 receptor.

- 5 In the present invention there is provided a compound of general formula (I)



I

wherein

- 10 R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently hydrogen, halo, C<sub>1</sub>-C<sub>6</sub> alkyl, -O(C<sub>1</sub>-C<sub>6</sub> alkyl), -CON(R<sup>9</sup>)<sub>2</sub>, -SOR<sup>9</sup>, -SO<sub>2</sub>R<sup>9</sup>, -SO<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, -N(R<sup>9</sup>)<sub>2</sub>, -NR<sup>9</sup>COR<sup>9</sup>, -CO<sub>2</sub>R<sup>9</sup>, -COR<sup>9</sup>, -SR<sup>9</sup>, -OH, -NO<sub>2</sub> or -CN;

each R<sup>9</sup> is independently hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl;

- R<sup>5</sup> and R<sup>6</sup> are each independently hydrogen, or C<sub>1</sub>-C<sub>6</sub> alkyl or together with the  
15 carbon atom to which they are attached form a C<sub>3</sub>-C<sub>7</sub> cycloalkyl group;

R<sup>7</sup> is hydrogen or C<sub>1</sub>-C<sub>6</sub> alkyl

n is 1 or 2;

X is a bond or, when n is 2, X may also be a NR<sup>9</sup> group;

wherein R<sup>9</sup> is as defined above;

- 20 when X is a bond R<sup>8</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, biphenyl or a 9-14 membered bicyclic or tricyclic heteroaryl group;

when X is a NR<sup>9</sup> group R<sup>8</sup> may additionally be phenyl, naphthyl or a 5-7 membered heteroaromatic ring; and

- 25 the R<sup>8</sup> group is optionally substituted with one or more substituents selected from

halo, C<sub>1</sub>-C<sub>6</sub> alkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, -O-aryl, heteroaryl, -O-heteroaryl, -CON(R<sup>9</sup>)<sub>2</sub>, -SOR<sup>9</sup>, -SO<sub>2</sub>R<sup>9</sup>, SO<sub>2</sub>N(R<sup>9</sup>)<sub>2</sub>, -N(R<sup>9</sup>)<sub>2</sub>, -NR<sup>9</sup>COR<sup>9</sup>, -CO<sub>2</sub>R<sup>9</sup>, -COR<sup>9</sup>, -SR<sup>9</sup>, -OH, -NO<sub>2</sub> or -CN;

wherein R<sup>9</sup> is as defined above;

5

or a pharmaceutically acceptable salt, hydrate, solvate, complex or prodrug thereof.

The compounds of general formula (I) are antagonists of PGD<sub>2</sub> at the CRTH2 receptor and will therefore be useful in the treatment of conditions which are mediated by PGD<sub>2</sub> binding to CRTH2. These include allergic diseases, asthmatic conditions and inflammatory diseases, examples of which are allergic asthma, perennial allergic rhinitis, seasonal allergic rhinitis, atopic dermatitis, contact hypersensitivity (including contact dermatitis), conjunctivitis, especially allergic conjunctivitis, eosinophilic bronchitis, food allergies, eosinophilic gastroenteritis, inflammatory bowel disease, ulcerative colitis and Crohn's disease, mastocytosis and also other PGD<sub>2</sub>-mediated diseases, for example autoimmune diseases such as hyper IgE syndrome and systemic lupus erythematus, psoriasis, acne, multiple sclerosis, allograft rejection, reperfusion injury, chronic obstructive pulmonary disease, as well as rheumatoid arthritis, psoriatic arthritis and osteoarthritis.

20

Similar, but not identical, compounds are disclosed in WO-A-9950268. These compounds differ from those of the present invention in that they do not contain a sulfone/sulfonamide moiety attached to the 3-position of the indole ring. In addition, they are not taught to be useful in the treatment of conditions such as asthma and allergic conditions, which are mediated by PGD<sub>2</sub>. Rather, they are said to be of use in the treatment of complications arising from diabetes mellitus.

25

PL 65781 and JP 43-24418 also relate to indole derivatives. However, the compounds disclosed in both of these documents differ from the compounds of the present application in that they are indole N-sulfonamides rather than 3-sulfones or 3-sulfonamides like the compounds of the present invention. The compounds

30

disclosed in PL 65781 and JP 43-24418 are similar in structure to indomethacin and, like indomethacin, are said to have anti-inflammatory and antipyretic activity. Thus, although this may not have been appreciated at the time when these documents were published, the compounds they describe are COX inhibitors, an activity which is quite different from that of the compounds of the present invention. Indeed, COX inhibitors are contraindicated in the treatment of many of the diseases and conditions, for example asthma and inflammatory bowel disease, for which the compounds of the present invention are useful, although they may sometimes be used to treat arthritic conditions.

10

Compounds which bind to the CRTH2 receptor are disclosed in WO-A-03/097042 and WO-A-03/097598. These compounds are indole acetic acids but in WO-A-03/097042 the indole system is fused at the 2-3 positions to a 5-7 membered carbocyclic ring. In WO-A-03/097598 there is a pyrrolidine group at the indole 3-position.

15

WO-A-03/101981 and WO-A-03/101961 both relate to CRTH2 antagonists. The compounds described in WO-A-03/101961 are similar in structure to the compounds of the present invention in which X is a bond. They differ from the compounds of general formula (I) because there is an -S- group linked to the indole 3-position in place of the SO or SO<sub>2</sub> group of the compounds of general formula (I). In addition, the group equivalent to the R<sup>8</sup> group in the compounds of general formula (I) is an aryl or heteroaryl group. There are no aliphatic substituents at this position as with the compounds of general formula (I). It has surprisingly been found that although these compounds have high intrinsic activity, they are less suitable for use as medicaments than the compounds of the present invention. This is because certain of the compounds of WO-A-03/101961 are inhibitors of cytochrome P<sub>450</sub>s and this has implications for the metabolism of any pharmacological agent which may be co-administered with these compounds. In contrast, the present inventors have shown that, surprisingly, the compounds of the present invention do not inhibit cytochrome P<sub>450</sub>s. In addition, our preliminary binding experiments have indicated that the

20

25

30

sulfide compounds described in WO-A-03/101961 appear to bind human eosinophils with a low off rate, which could lead to an unpredictable duration of action.

WO-A-03/10981 relates to compounds which are of similar structure to the compounds of the present invention except that the substituent at the 3-position of the indole ring system is a phenyl, naphthyl or heteroaryl group with no SO, SO<sub>2</sub> or SO<sub>2</sub>NR<sup>9</sup> linker as with the compounds of general formula (I). Clearly, the inclusion of a linking group is likely to have a substantial effect on the activity of the compound. Furthermore, the substituent at the indole 3-position cannot be an aliphatic group as in the present invention.

WO-A-2004/007451 relates to CRTH2 inhibitors which are similar in structure to the compounds of the present invention in which X is a bond, except that the group equivalent to the R<sup>8</sup> group of the compounds of general formula (I) is phenyl, naphthyl or a 5-7 membered heteroaromatic group. In fact, all the exemplified compounds have a substituted phenyl group at this position. This is clearly different from the compounds of the present invention where the R<sup>8</sup> groups are either a bicyclic or tricyclic heteroaromatic ring or an alkyl, alkenyl or alkynyl group. It is particularly surprising that compounds containing alkyl, alkenyl and alkynyl groups have proved to be so active since they differ markedly in structure from the prior art compounds.

In the present specification "C<sub>1</sub>-C<sub>6</sub> alkyl" refers to a straight or branched saturated hydrocarbon chain having one to six carbon atoms and optionally substituted with one or more halo substituents or with one or more C<sub>3</sub>-C<sub>7</sub> cycloalkyl groups. Examples include methyl, ethyl, n-propyl, isopropyl, t-butyl, n-hexyl, trifluoromethyl, 2-chloroethyl, methylenecyclopropyl, methylenecyclobutyl, methylenecyclobutyl and methylenecyclopentyl.

"C<sub>1</sub>-C<sub>4</sub> alkyl" and "C<sub>1</sub>-C<sub>18</sub> alkyl" have similar meanings except that they contain from one to four and from one to eighteen carbon atoms respectively.

C<sub>3</sub>-C<sub>7</sub> cycloalkyl refers to a saturated 3 to 7 membered carbocyclic ring. Examples of such groups include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

5 The terms "C<sub>2</sub>-C<sub>6</sub> alkenyl" and "C<sub>2</sub>-C<sub>6</sub> alkynyl" refer straight or branched hydrocarbon chains having from two to six carbon atoms and containing respectively at least one carbon-carbon double bond or at least one carbon-carbon triple bond. As with alkyl groups they may optionally be substituted with one or more halo substituents or with one or more C<sub>3</sub>-C<sub>7</sub> cycloalkyl groups.

10 In the present specification, "halo" refers to fluoro, chloro, bromo or iodo.

The terms "aromatic moiety" and "aryl" in the context of the present specification refer to an aromatic ring system having from 5 to 14 ring carbon atoms and containing up to three rings. Examples of aromatic moieties are benzene and  
15 naphthalene. Aryl groups may be substituted with one or more substituents chosen from halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, a 5-7-membered heterocyclic ring or SO<sub>2</sub>R<sup>9</sup> where R<sup>9</sup> is as defined above.

The terms "heteroaromatic moiety" and "heteroaryl" refer to an aromatic ring system  
20 in which at least one of the ring carbon atoms is replaced by a nitrogen, oxygen or sulfur atom. Examples include single ring systems such as pyridine, pyrimidine, pyrazole, thiophene, oxazole and isoxazole. Other examples include fused ring systems such as quinoline, isoquinoline, quinazoline, benzthiazole, benzoxazole, benzimidazole and indole groups.

25

Unless stated otherwise a heteroaromatic moiety has from 5 to 14 ring carbon atoms but, for example, "5-7 membered heteroaromatic ring" contains 5 to 7 ring atoms. Bicyclic and tricyclic heteroaryl groups contain respectively two or three fused rings. Bicyclic heteroaryl groups may be, for example, 6,6- or 6-5-ring systems such as  
30 those exemplified above.



As with aryl groups, heteroaryl groups may also be substituted with one or more substituents chosen from halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, a 5-7-membered heterocyclic ring or SO<sub>2</sub>R<sup>9</sup> where R<sup>9</sup> is as defined above.

- 5     The term "5 to 7 membered heterocyclic ring" refers to a non-aromatic ring system having from 5 to 7 ring atoms and wherein at least one of the ring carbon atoms is replaced by a nitrogen, oxygen or sulfur atom. Examples include piperidine, morpholine, imidazoline, piperazine and tetrahydrofuran.
- 10    Appropriate pharmaceutically and veterinarily acceptable salts of the compounds of general formulae (I) and (II) include basic addition salts such as sodium, potassium, calcium, aluminium, zinc, magnesium and other metal salts as well as choline, diethanolamine, ethanolamine, ethyl diamine and other well known basic addition salts.
- 15     Where appropriate, pharmaceutically or veterinarily acceptable salts may also include salts of organic acids, especially carboxylic acids, including but not limited to acetate, trifluoroacetate, lactate, gluconate, citrate, tartrate, maleate, malate, pantothenate, adipate, alginate, aspartate, benzoate, butyrate, digluconate,
- 20    cyclopentanate, glucoheptanate, glycerophosphate, oxalate, heptanoate, hexanoate, fumarate, nicotinate, pamoate, pectinate, 3-phenylpropionate, picrate, pivalate, propionate, tartrate, lactobionate, pivate, camphorate, undecanoate and succinate, organic sulfonic acids such as methanesulfonate, ethanesulfonate, 2-hydroxyethane sulfonate, camphorsulfonate, 2-naphthalenesulfonate, benzenesulfonate, p-
- 25    chlorobenzenesulfonate and p-toluenesulfonate; and inorganic acids such as hydrochloride, hydrobromide, hydroiodide, sulfate, bisulfate, hemisulfate, thiocyanate, persulfate, phosphoric and sulfonic acids.

Salts which are not pharmaceutically or veterinarily acceptable may still be valuable

30    as intermediates.

Prodrugs are any covalently bonded compounds which release the active parent drug according to general formula (I) *in vivo*. Examples of prodrugs include alkyl esters of the compounds of general formula (I), for example the esters of general formula (II) below.

5

If a chiral centre or another form of isomeric centre is present in a compound of the present invention, all forms of such isomer or isomers, including enantiomers and diastereoisomers, are intended to be covered herein. Compounds of the invention containing a chiral centre may be used as a racemic mixture, an enantiomerically enriched mixture, or the racemic mixture may be separated using well-known techniques and an individual enantiomer may be used alone.

10

In the compounds of general formula (I), it is preferred that, independently or in any combination:

15

$R^1$  is halo or hydrogen;

$R^2$  is halo or hydrogen;

$R^3$  is halo or hydrogen;

$R^4$  is halo or hydrogen.

20

In more preferred compounds,  $R^1$ ,  $R^3$  and  $R^4$  are hydrogen, while  $R^2$  is halo, particularly fluoro.

In preferred compounds of general formula (I),  $R^5$  and  $R^6$  are each independently hydrogen or  $C_1$ - $C_4$  alkyl. However, in more active compounds, at least one, and preferably both of  $R^5$  and  $R^6$  are hydrogen.

25

Compounds of general formula (I) preferably have an  $R^7$  group chosen from H or  $C_1$ - $C_6$  alkyl; most suitably  $R^7$  is methyl.

30

In particularly preferred compounds of general formula (I), n is 2.

When X is a bond, it is preferred that  $R^8$  is  $C_1$ - $C_6$  alkyl, biphenyl or a bicyclic heteroaryl group, any of which may be substituted with halogen, phenyl,  $-CO_2R^9$ ,  $CON(R^9)_2$  or  $-SO_2R^9$ , where  $R^9$  is as defined above.

- 5 More preferred compounds in which X is a bond include those in which  $R^8$  is  $C_1$ - $C_4$  alkyl, biphenyl or a bicyclic heteroaryl group, any of which may be substituted with phenyl,  $-CO_2R^9$ ,  $CON(R^9)_2$  or  $-SO_2R^9$ , where  $R^9$  is H or  $C_1$ - $C_4$  alkyl.

When X is  $NR^9$ , it is preferred that  $R^9$  is H or methyl and  $R^8$  is:


- 10 phenyl optionally substituted with one or more halo,  $C_1$ - $C_6$ alkyl or  $-O(C_1$ - $C_6$  alkyl) groups;  
 $C_1$ - $C_6$  alkyl, optionally substituted with aryl; or  
heteroaryl.
- 15 More preferably, when X is  $NR^9$ ,  $R^8$  is phenyl, benzyl or pyridyl, any of which may optionally be substituted with one or more halo, methyl or methoxy groups.

Among the most preferred compounds are the following:

1. [3-(Butane-1-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 20 2. 3-(Biphenyl-4-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
3. (3-Carboxymethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid
4. (3-Carbamoylmethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid
5. [5-Fluoro-3-(2-methanesulfonyl-ethanesulfonyl)-2-methyl-indol-1-yl]-acetic acid
6. [3-(Benzothiazole-2-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 25 7. [3-(Benzothiazole-2-sulfinyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
8. [5-Fluoro-2-methyl-3-(quinoline-2-sulfonyl)-indol-1-yl]-acetic acid
9. [5-Fluoro-2-methyl-3-(quinolin-8-ylsulfonyl)-indol-1-yl]-acetic acid
10. (5-Fluoro-2-methyl-3-phenylmethanesulfonyl-1H-indol-1-yl)-acetic acid
11. [3-(4-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 30 12. [3-(3-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
13. [3-(4-Fluoro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid

14. [3-(2-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
 15. (3-Benzylsulfamoyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid  
 16. [5-Fluoro-3-(2-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid  
 17. [5-Fluoro-3-(4-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid  
 5 18. (5-Fluoro-2-methyl-3-phenylsulfamoyl-indol-1-yl)-acetic acid  
 19. [3-(3,4-Dichloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
 20. [5-Fluoro-3-(3-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid  
 21. (5-Fluoro-2-methyl-3-*m*-tolylsulfamoyl-indol-1-yl)-acetic acid  
 22. (5-Fluoro-2-methyl-3-*p*-tolylsulfamoyl-indol-1-yl)-acetic acid  
 10 23. [3-(4-Chloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
 24. [3-(Benzyl-methyl-sulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
 25. [5-Fluoro-2-methyl-3-(pyridin-3-ylsulfamoyl)-indol-1-yl]-acetic acid;  
 or the C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, (CH<sub>2</sub>)<sub>m</sub>OC(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, (CH<sub>2</sub>)<sub>m</sub>N(R<sup>11</sup>)<sub>2</sub>,  
 CH((CH<sub>2</sub>)<sub>m</sub>O(C=O)R<sup>12</sup>)<sub>2</sub> esters of any of the above; wherein  
 15 m is 1 or 2;  
 R<sup>11</sup> is hydrogen or methyl;  
 R<sup>12</sup> is C<sub>1</sub>-C<sub>18</sub> alkyl.

In a further aspect of the present invention, there is provided a compound of general  
 20 formula (II):



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, n, X, R<sup>7</sup> and R<sup>8</sup> are as defined for general formula (I);  
 R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, (CH<sub>2</sub>)<sub>m</sub>OC(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, (CH<sub>2</sub>)<sub>m</sub>N(R<sup>11</sup>)<sub>2</sub>,  
 25 CH((CH<sub>2</sub>)<sub>m</sub>O(C=O)R<sup>12</sup>)<sub>2</sub>;

m is 1 or 2;


R<sup>11</sup> is hydrogen or methyl;

R<sup>12</sup> is C<sub>1</sub>-C<sub>18</sub> alkyl.

- 5 Compounds of general formula (II) are novel and may be used as prodrugs for compounds of general formula (I). When the compound of general formula (II) acts as a prodrug, it is later transformed to the drug by the action of an esterase in the blood or in a tissue of the patient.
- 10 Examples of particularly suitable R<sup>10</sup> groups when the compound of general formula (II) is used as a prodrug include:
- methyl, ethyl, propyl, phenyl, CH<sub>2</sub>OC(=O)tBu, CH<sub>2</sub>CH<sub>2</sub>N(Me)<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub> or CH(CH<sub>2</sub>O(C=O)R<sup>12</sup>)<sub>2</sub> wherein R<sup>12</sup> is as defined above.
- 15 Compounds of general formula (I) wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are as defined for general formula (I) and X is a bond, may be prepared from compounds of general formula (Ia), which is a compound of general formula (I) wherein n is 0 and X is a bond, by oxidation with a suitable oxidising agent such as potassium peroxymonosulfate, m-CPBA, hydrogen peroxide or other well known oxidising
- 20 reagents.

- In addition to their use as prodrugs, compounds of formula (II) wherein R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl may be used in a process for the preparation of a compound of general formula (I), the process comprising reacting the compound of general formula (II) with a base
- 25 such as sodium hydroxide or lithium hydroxide. The reaction may take place in an aqueous solvent or an organic solvent or a mixture of the two. A typical solvent used for the reaction is a mixture of tetrahydrofuran and water. The same method may be used to prepare compounds of general formula (Ia) as defined above from compounds of general formula (IIa), which are identical to compounds of general
- 30 formula (II) except that n is 0.

Compounds of general formula (II) and (IIa) in which X is a bond may be prepared from compounds of general formula (III):



III

- 5 wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^7$  and  $R^8$  are as defined for general formula (I) and  $n$  is 0, 1 or 2;

by reaction with a compound of general formula (IV):



10

wherein  $R^5$  and  $R^6$  are as defined for general formula (I),  $R^{10}$  is as defined for general formula (II) and X is a leaving group in particular a halo group, for example bromo.


The reaction is conducted under strongly basic conditions, for example in the presence of excess sodium hydride, and in a polar organic solvent such as dimethylformamide.

15

Compounds of general formula (IV) are well known and are readily available or can be prepared by methods known to those skilled in the art.

- 20 Compounds of general formula (III) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^7$  and  $R^8$  are as defined for general formula (I) and  $n$  is 2 can be prepared by reacting a compound of general formula (V):

14



wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^7$  are as defined in general formula (I);

5

with a compound of general formula (VI):




10 wherein  $R^8$  is as defined in general formula (I).

The reaction is carried out in the presence of a Lewis acid such as indium(III) bromide. The reaction may be conducted in a polar organic solvent, particularly a chlorinated solvent such as 1,2-dichloroethane

15

Compounds of general formulae (V) and (VI) are well known in the art and are readily available or can be prepared by known methods.

Compounds of general formula (II) in which X is  $NR^9$  may be prepared from compounds of general formula (VII):



20

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$  and  $R^7$  are as defined for general formula (I) and  $R^{10}$  is

as defined in general formula (II) by reaction with a compound of general formula (VIII):




5

wherein  $\text{R}^8$  and  $\text{R}^9$  is as defined above for general formula (I).

The reaction solvent may be a polar organic solvent such as dichloromethane.

10 Compounds of general formulae (VIII) are well known and are either readily available or can be prepared by methods well known to those skilled in the art.

Compounds of general formula (VII) may be prepared from compounds of general formula (IX)



15

IX

wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$ , and  $\text{R}^7$  are as defined in general formula (I) and  $\text{R}^{10}$  is as defined for general formula (II);

by reaction with chlorosulfonic acid.

20

The reaction preferably takes place in a non polar organic solvent.

Compounds of general formula (IX) are well known and are readily available or can be prepared by methods well known to those skilled in the art.

25

Compounds of general formula (III) wherein  $\text{R}^1$ ,  $\text{R}^2$ ,  $\text{R}^3$ ,  $\text{R}^4$ ,  $\text{R}^7$  and  $\text{R}^8$  are as defined



for general formula (I) and n is 0 can be prepared by reacting a compound of general formula (IX) wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^7$  are as defined in general formula (I) and  $R^{10}$  is as defined for general formula (II) with a compound of general formula (X):



wherein  $R^8$  is as defined in general formula (I).

10            The reaction is carried out in the presence of iodine and potassium iodide. The reaction may take place in an aqueous or an organic solvent or a mixture of the two. A typical solvent used for the reaction is a mixture such as ethanol and water.

15            Compounds of general formula (I) are antagonists of  $PGD_2$  at the CRTH2 receptor and compounds of general formula (II) are prodrugs for compounds of general formula (I). Compounds of general formulae (I) and (II) are therefore useful in a method for the treatment of diseases and conditions mediated by  $PGD_2$  at the CRTH2 receptor, the method comprising administering to a patient in need of such treatment a suitable amount of a compound of general formula (I) or (II).

20            In a third aspect of the invention, there is provided a compound of general formula (I) or (II) for use in medicine, particularly for use in the treatment or prevention of diseases and conditions mediated by  $PGD_2$  at the CRTH2 receptor.

25            Furthermore, there is also provided the use of a compound of general formula (I) or (II) in the preparation of an agent for the treatment or prevention of diseases and conditions mediated by  $PGD_2$  at the CRTH2 receptor.

30            As mentioned above, such diseases and conditions include allergic asthma, perennial allergic rhinitis, seasonal allergic rhinitis, atopic dermatitis, contact hypersensitivity (including contact dermatitis), conjunctivitis, especially allergic conjunctivitis, eosinophilic bronchitis, food allergies, eosinophilic gastroenteritis, inflammatory

bowel disease, ulcerative colitis and Crohn's disease, mastocytosis and also other PGD<sub>2</sub>-mediated diseases, for example autoimmune diseases such as hyper IgE syndrome and systemic lupus erythematus, psoriasis, acne, multiple sclerosis, allograft rejection, reperfusion injury, chronic obstructive pulmonary disease, as well  
5 as rheumatoid arthritis, psoriatic arthritis and osteoarthritis.

The compounds of general formula (I) or (II) must be formulated in an appropriate manner depending upon the diseases or conditions they are required to treat.

10 Therefore, in a further aspect of the invention there is provided a pharmaceutical composition comprising a compound of general formula (I) or (II) together with a pharmaceutical excipient or carrier. Other active materials may also be present, as may be considered appropriate or advisable for the disease or condition being treated or prevented.

15

The carrier, or, if more than one be present, each of the carriers, must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient.

20 The formulations include those suitable for oral, rectal, nasal, bronchial (inhaled), topical (including eye drops, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration and may be prepared by any methods well known in the art of pharmacy.

25 The route of administration will depend upon the condition to be treated but preferred compositions are formulated for oral, nasal, bronchial or topical administration.

The composition may be prepared by bringing into association the above defined  
30 active agent with the carrier. In general, the formulations are prepared by uniformly and intimately bringing into association the active agent with liquid carriers or finely

divided solid carriers or both, and then if necessary shaping the product. The invention extends to methods for preparing a pharmaceutical composition comprising bringing a compound of general formula (I) or (II) in conjunction or association with a pharmaceutically or veterinarily acceptable carrier or vehicle.

5

Formulations for oral administration in the present invention may be presented as: discrete units such as capsules, sachets or tablets each containing a predetermined amount of the active agent; as a powder or granules; as a solution or a suspension of the active agent in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water  
10 liquid emulsion or a water in oil liquid emulsion; or as a bolus etc.

For compositions for oral administration (e.g. tablets and capsules), the term “acceptable carrier” includes vehicles such as common excipients e.g. binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, polyvinylpyrrolidone  
15 (Povidone), methylcellulose, ethylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, sucrose and starch; fillers and carriers, for example corn starch, gelatin, lactose, sucrose, microcrystalline cellulose, kaolin, mannitol, dicalcium phosphate, sodium chloride and alginic acid; and lubricants such as magnesium stearate, sodium stearate and other metallic stearates, glycerol stearate  
20 stearic acid, silicone fluid, talc waxes, oils and colloidal silica. Flavouring agents such as peppermint, oil of wintergreen, cherry flavouring and the like can also be used. It may be desirable to add a colouring agent to make the dosage form readily identifiable. Tablets may also be coated by methods well known in the art.

25 A tablet may be made by compression or moulding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active agent in a free flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, preservative, surface-active or dispersing agent. Moulded tablets may be made by moulding in a  
30 suitable machine a mixture of the powdered compound moistened with an inert liquid diluent. The tablets may optionally be coated or scored and may be formulated

so as to provide slow or controlled release of the active agent.

Other formulations suitable for oral administration include lozenges comprising the active agent in a flavoured base, usually sucrose and acacia or tragacanth; pastilles  
5 comprising the active agent in an inert base such as gelatin and glycerin, or sucrose and acacia; and mouthwashes comprising the active agent in a suitable liquid carrier.

For topical application to the skin, compounds of general formula (I) or (II) may be made up into a cream, ointment, jelly, solution or suspension etc. Cream or ointment  
10 formulations that may be used for the drug are conventional formulations well known in the art, for example, as described in standard text books of pharmaceutics such as the British Pharmacopoeia.

Compounds of general formula (I) or (II) may be used for the treatment of the  
15 respiratory tract by nasal, bronchial or buccal administration of, for example, aerosols or sprays which can disperse the pharmacological active ingredient in the form of a powder or in the form of drops of a solution or suspension. Pharmaceutical compositions with powder-dispersing properties usually contain, in addition to the active ingredient, a liquid propellant with a boiling point below room temperature  
20 and, if desired, adjuncts, such as liquid or solid non-ionic or anionic surfactants and/or diluents. Pharmaceutical compositions in which the pharmacological active ingredient is in solution contain, in addition to this, a suitable propellant, and furthermore, if necessary, an additional solvent and/or a stabiliser. Instead of the propellant, compressed air can also be used, it being possible for this to be produced  
25 as required by means of a suitable compression and expansion device.

Parenteral formulations will generally be sterile.

Typically, the dose of the compound will be about 0.01 to 100 mg/kg; so as to  
30 maintain the concentration of drug in the plasma at a concentration effective to inhibit PGD<sub>2</sub> at the CRTH2 receptor. The precise amount of a compound of general

formula (I) or (II) which is therapeutically effective, and the route by which such compound is best administered, is readily determined by one of ordinary skill in the art by comparing the blood level of the agent to the concentration required to have a therapeutic effect.

5

Compounds of general formula (I) or (II) may be used in combination with one or more active agents which are useful in the treatment of the diseases and conditions listed above, although these active agents are not necessarily inhibitors of PGD<sub>2</sub> at the CRTH2 receptor.

10

Therefore, the pharmaceutical composition described above may additionally contain one or more of these active agents.

There is also provided the use of a compound of general formula (I) or (II) in the preparation of an agent for the treatment of diseases and conditions mediated by PGD<sub>2</sub> at the CRTH2 receptor, wherein the agent also comprises an additional active agent useful for the treatment of the same diseases and conditions.

15

These additional active agents which may have a completely different mode of action include existing therapies for allergic and other inflammatory diseases including:

20

β<sub>2</sub> agonists such as salmeterol;

corticosteroids such as fluticasone;

antihistamines such as loratidine;

leukotriene antagonists such as montelukast;

25

anti-IgE antibody therapies such as omalizumab;

anti-infectives such as fusidic acid (particularly for the treatment of atopic dermatitis);

anti-fungals such as clotrimazole (particularly for the treatment of atopic dermatitis);

immunosuppressants such as tacrolimus and particularly pimecrolimus in the case of

30

inflammatory skin disease.

CRTH2 antagonists may also be combined with therapies that are in development for

inflammatory indications including:

other antagonists of PGD<sub>2</sub> acting at other receptors such as DP antagonists;

inhibitors of phosphodiesterase type 4 such as cilonilast;

drugs that modulate cytokine production such as inhibitors of TNF $\alpha$  converting

5 enzyme (TACE);

drugs that modulate the activity of Th2 cytokines IL-4 and IL-5 such as blocking monoclonal antibodies and soluble receptors;

PPAR- $\gamma$  agonists such as rosiglitazone;

5-lipoxygenase inhibitors such as zileuton.

10

In yet a further aspect of the invention, there is provided a product comprising a compound of general formula (I) or (II) and one or more of the agents listed above as a combined preparation for simultaneous, separate or sequential use in the treatment of a disease or condition mediated by the action of PGD<sub>2</sub> at the CRTH2 receptor.

15

The invention will now be described in greater detail with reference to the following non limiting examples and the drawings in which:

Figure 1 shows the effects of CRTH2 agonists on calcium mobilisation in  
20 CHO/CRTH2 cells.

### **Example 1 - Synthesis of 3-Sulfonyl indole Derivatives (Method A)**

#### **1. Synthesis of 3-(Butane-1-sulfonyl)-5-fluoro-2-methyl-1H-indole**

25 Indium (III) bromide (94.7 mg, 0.267 mmol) was added in one portion to a stirred solution of 5-fluoro-2-methylindole (50 mg, 0.34 mmol) and butanesulfonyl chloride (418 mg, 2.67 mmol) in 1,2-dichloroethane (2 ml) at room temperature. The mixture was subjected to microwave conditions (85 °C, 150 W) for 45 minutes, cooled to room temperature and then concentrated *in vacuo* to leave a brown residue.  
30 Purification by flash column chromatography on silica gel eluting with 10 % ethyl acetate : hexane to 100 % ethyl acetate gave the *sulfone* (55 mg, 15 %) as an off-

white solid.

2. **Synthesis of [3-(Butane-1-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid (Compound 1)**

3-(Butane-1-sulfonyl)-5-fluoro-2-methyl-1H-indole (55 mg, 0.204 mmol) in DMF (1 ml) was added dropwise over 1 minute to a stirred suspension of sodium hydride (11 mg, 0.29 mmol; 60 % in mineral oil) in DMF (1 ml) at 0 °C. The solution was stirred at 0 °C for 45 minutes and then ethyl bromoacetate (0.032 ml, 0.29 mmol) was added dropwise and the resulting mixture stirred at room temperature for 18 hours. The mixture was adjusted to pH 4 with 10 % citric acid and then extracted into ethyl acetate (2 x 10 ml). The combined organic extracts were dried and concentrated *in vacuo* to leave a residue. The residue was taken up into THF (1 ml) and lithium hydroxide monohydrate (19 mg, 0.464 mmol) in water (1 ml) was then added in one portion at room temperature. The mixture was stirred at room temperature for 3 hours and then the solution adjusted to pH 4 with 10 % citric acid. The product was extracted with ethyl acetate and the combined organic extracts were dried and concentrated *in vacuo* to leave a residue which was triturated with diethyl ether to give the *carboxylic acid* as an off-white solid (5.4 mg, 8 %),  $\delta_H$  (400 MHz, MeOD) 7.57 (1H, dd  $J$  9.8, 2.3 Hz, *Ar*), 7.43 (1H, dd  $J$  9.1, 4.0 Hz, *Ar*), 7.04 (1H, td  $J$  9.1, 2.5 Hz, *Ar*), 4.79 (2H, s,  $CH_2CO_2H$ ), 3.23-3.19 (2H, m,  $SO_2CH_2$ ), 2.70 (3H, s,  $CCH_3$ ), 1.77-1.70 (2H, m,  $CH_2CH_2CH_2CH_3$ ), 1.47-1.41 (2H, m,  $CH_2CH_2CH_2CH_3$ ), 0.93 (3H, t  $J$  7.6 Hz,  $CH_2CH_2CH_2CH_3$ ); Tr = 1.38 min,  $m/z$  ( $ES^+$ ) ( $M+H$ )<sup>+</sup> 308.24. Tr = 1.82 min (98 %),  $m/z$  ( $ES^+$ ) ( $M+H$ )<sup>+</sup> 328.20.

Compound 2 was prepared using the same general method as for Compound 1 but with appropriately chosen starting materials.

**Compound 2 – 3-(Biphenyl-4-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**  
 $\delta_H$  (400 MHz, MeOD) 8.03 (2H, d,  $J$  8.6 Hz *Ar*), 7.80 (2H, d,  $J$  8.6 Hz, *Ar*), 7.77-7.74 (1H, dd,  $J$  9.6, 2.5 Hz, *Ar*), 7.66-7.64 (2H, dd,  $J$  8.0, 1.3 Hz, *Ar*), 7.49-7.39 (4H,

m, Ar), 7.07 (1H, td,  $J$  9.1, 2.5 Hz, Ar), 5.07 (2H, s,  $\text{CH}_2$ ), 2.76 (3H, s,  $\text{CH}_3$ ); Tr = 1.52 min,  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 424.1.

**Example 2 - Synthesis of 3-Sulfonyl indole Derivatives (Method B)**

5

**1. 2-Methylsulfanyl-ethanethiol**

A solution of methyl iodide (10 ml, 22.82 g, 0.161 mol) in acetone (50ml) was added dropwise to a stirred suspension of ethane dithiol (11.24 ml, 12.62 g, 0.134 mol) and potassium carbonate (37.04 g, 0.268 mol) in acetone (150 ml). The resulting mixture was stirred at room temperature for 4 hours. Water (150 ml) was added, and the mixture stirred for a further 15 minutes. The reaction mixture was extracted with dichloromethane (3 x 200 ml), the organic washings combined, dried over sodium sulfate and evaporated (maintaining pressure above 200 mbar to ensure no co-evaporation of product) to give 2-methylsulfanyl-ethanethiol. LC/MS showed <5% starting material and a 2:1 mixture of mono and bis-methylated material. The material was used in the next step with no further purification.

**2. [5-Fluoro-2-methyl-3-(2-methylsulfanyl-ethylsulfanyl)-indol-1-yl]-acetic acid ethyl ester**

To a stirred solution of (5-fluoro-2-methyl-indol-1-yl)-acetic acid ethyl ester (1.20g, 5.10mmol) and 2-methylsulfanyl-ethanethiol (1.04g, 6.12mmol) in 1:1 EtOH:H<sub>2</sub>O (40ml) at room temperature was added iodine (1.29g, 5.10mmol) and potassium iodide (0.847g, 5.10mmol). The mixture was heated to 100°C and stirred for 2 hours, then stirred at room temperature for 16 hours. The reaction mixture was quenched by the careful addition of saturated NaHCO<sub>3</sub> (aq), then extracted with DCM (2 x 50ml). The organic washings were combined, washed with saturated sodium thiosulfate (2 x 70ml), dried over magnesium sulfate and evaporated to give an oil. The crude oil was purified by chromatography (3 x 12cm column; 4:1 hexane:EtOAc as eluant) to give the ester (1.17g, 67%).  $\delta_{\text{H}}$  (400 MHz, CDCl<sub>3</sub>) 7.36 (1H, dd  $J$  9.2, 2.5 Hz, Ar), 7.11 (1H, dd  $J$  8.8, 4.1 Hz, Ar); 7.05 (1H, td  $J$  9.2, 2.3 Hz, Ar), 4.79 (2H, s,  $\text{CH}_2\text{CO}_2\text{Et}$ ), 4.21 (2H, q  $J$  7.2 Hz,  $\text{CO}_2\text{CH}_2\text{CH}_3$ ), 2.86-2.79 (2H, m,



$\text{CH}_2\text{CH}_2$ ), 2.59-2.55 (2H, m,  $\text{CH}_2\text{CH}_2$ ), 2.50 (3H, s,  $\text{SCH}_3$ ), 2.04 (3H, s,  $\text{CCH}_3$ ), 1.25 (3H, t J 7.2 Hz,  $\text{CO}_2\text{CH}_2\text{CH}_3$ ).

3. **[5-Fluoro-3-(2-methanesulfonyl-ethanesulfonyl)-2-methyl-indol-1-yl]-acetic acid ethyl ester**

Oxone (8.43g, 13.7mmol) was added to a stirred solution of [5-fluoro-2-methyl-3-(2-methylsulfonyl-ethylsulfonyl)-indol-1-yl]-acetic acid ethyl ester (1.17g, 3.43mmol) in 4:1 1,4-dioxane: $\text{H}_2\text{O}$  at room temperature. After 30 minutes the reaction mixture was quenched by the careful addition of saturated sodium bicarbonate (50ml; care – effervescence), then extracted with DCM (2 x 100ml). The organic washings were combined and washed with brine (2 x 100ml). Aqueous washings were then back-extracted with DCM (100ml). All organic layers were combined, dried over magnesium sulfate and evaporated to give a pale green crystalline solid. The solid was suspended in DCM and collected *via* filtration to give the *ester* (1.06g, 76%).  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 7.66 (1H, dd J 9.1, 2.5 Hz, Ar), 7.22 (1H, dd J 9.0, 4.0 Hz, Ar), 7.08 (1H, td J 8.9, 2.5 Hz, Ar), 4.86 (2H, s,  $\text{CH}_2\text{CO}_2\text{Et}$ ), 4.26 (2H, q J 7.1 Hz,  $\text{CO}_2\text{CH}_2\text{CH}_3$ ), 3.61-3.57 (2H, m,  $\text{CH}_2\text{CH}_2$ ), 3.48-3.44 (2H, m,  $\text{CH}_2\text{CH}_2$ ), 2.99 (3H, s,  $\text{SCH}_3$ ), 2.71 (3H, s,  $\text{CH}_3$ ), 1.30 (3H, t J 7.1 Hz,  $\text{CO}_2\text{CH}_2\text{CH}_3$ ).

4. **Compound 5 – [[5-Fluoro-3-(2-methanesulfonyl-ethanesulfonyl)-2-methyl-indol-1-yl]-acetic acid**

Lithium hydroxide monohydrate (132 mg, 3.14 mmol) was added in one portion to a stirred solution of [5-fluoro-3-(2-methanesulfonyl-ethanesulfonyl)-2-methyl-indol-1-yl]-acetic acid ethyl ester (1.06 g, 2.61 mmol) in THF : water (5:1; 15 ml) and the resulting mixture stirred at room temperature for 2 h. The mixture was concentrated *in vacuo* to leave a residue which was partitioned between ethyl acetate and 10 % citric acid. The organic layer was separated and the aqueous solution extracted with ethyl acetate (3 x 100 ml). The combined organic extracts were dried and concentrated *in vacuo* to leave an off-white solid. The solid was then triturated with dichloromethane to give the *carboxylic acid* as an off-white solid, (438 mg, 44 %),  $\delta_{\text{H}}$  (400 MHz,  $d_6$ -Acetone) 7.58 - 7.63 (2H, m, Ar) 7.09 (1H, td J 9.2, 2.6 Hz, Ar),

5.22 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 3.60-3.66 (2H, m,  $\text{SO}_2\text{CH}_2\text{CH}_2$ ), 3.45 (2H, m,  $\text{SO}_2\text{CH}_2\text{CH}_2$ ), 3.02 (3H, s,  $\text{SO}_2\text{CH}_3$ ), 2.75 (3H, s,  $\text{CH}_3$ ); Tr = 1.08 min (98 %),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 378.16.

Compounds 3 and 4 were prepared by a similar route using appropriate starting materials.

**Compound 3 – (3-Carboxymethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid**

Tr = 1.16 min,  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 330.10.

10

**Compound 4 – (3-Carbamoylmethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid**

Tr = 1.50 min,  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 329.17.

15 **Example 3 - Synthesis of 3-Sulfonyl indole Derivatives (Method B2)**

A similar method to that set out in step 2 of Example 2 above was used to synthesise the following intermediates. However, hydrolysis to the acid took place before oxidation to give the sulfone or sulfoxide derivative.

20

**[5-Fluoro-2-methyl-3-(quinolin-8-ylsulfanyl)-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz, MeOD) 8.95-8.94 (1H, m, Ar), 8.36 (1H, dd  $J$  8.3, 1.7 Hz, Ar), 7.64-7.60 (2H, m, Ar), 7.45 (1H, dd  $J$  8.8, 4.2 Hz, Ar), 7.29 (1H, t  $J$  7.8 Hz, Ar), 7.09 (1H, dd  $J$  9.2, 2.6 Hz, Ar), 7.00 (1H, td  $J$  9.2, 2.6 Hz, Ar), 6.85 (1H, app d  $J$  7.3 Hz, Ar), 5.14 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 2.52 (3H, s,  $\text{CCH}_3$ ); Tr = 1.30 min,  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 367.39.

25

**[5-Fluoro-2-methyl-3-(quinolin-2-ylsulfanyl)-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz, MeOD) 8.01 (1H, d  $J$  8.6 Hz, Ar), 7.93 (1H, d  $J$  7.8 Hz, Ar), 7.82 (1H, d  $J$  8.1 Hz, Ar), 7.76 (1H, app td  $J$  7.1, 1.4 Hz, Ar), 7.53 (1H, app td  $J$  7.0, 1.1 Hz, Ar), 7.47 (1H, dd  $J$  9.1, 4.2 Hz, Ar), 7.16 (1H, dd  $J$  9.0, 2.4 Hz, Ar), 7.03 (1H, td  $J$

30

9.2, 2.6 Hz, *Ar*), 6.87 (1H, d *J* 8.8 Hz, *Ar*), 5.14 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.55 (3H, s, CCH<sub>3</sub>); Tr = 1.37 min, *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 367.24.

**[3-(Benzothiazol-2-ylsulfanyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

5     δ<sub>H</sub> (400 MHz, MeOD) 7.81 (1H d *J* 8.3 Hz, *Ar*), 7.71 (1H, d *J* 7.8 Hz, *Ar*), 7.50-7.43 (2H, m, *Ar*), 7.31-7.24 (2H, m, *Ar*), 7.06 (1H td *J* 9.0, 2.4 Hz, *Ar*), 5.15 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.60 (3H, s, CCH<sub>3</sub>); Tr = 1.49 min, *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 373.34.

**(3-Benzylsulfanyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid**

10     δ<sub>H</sub> (250 MHz, *d*<sub>6</sub>-DMSO) 7.46 (1H, dd *J* 8.8, 4.3 Hz, *Ar*), 7.21-7.18 (3H, m, *Ar*), 7.14 (1H, dd *J* 9.5, 2.5 Hz, *Ar*), 7.01-6.92 (3H, m, *Ar*), 4.99 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 3.75 (2H, s, ArCH<sub>2</sub>), 2.01 (3H, s, CH<sub>3</sub>); Tr = 1.56min (100%) *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 330.16.

15     These intermediates can then be oxidised to give compounds of general formula (I) where n is 1 or 2 using the following method.

1.     **Compound 6 – [3-(Benzothiazole-2-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid and Compound 7 – [3-(Benzothiazole-2-sulfinyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

20

Potassium peroxymonosulfate (131.0 mg, 214 μmol) was added in one portion to a stirred solution of the [3-(benzothiazol-2-ylsulfanyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid, 20.0 mg, 53.6 μmol) in 1, 4-dioxane : water (0.3 ml; 4:1) at room temperature. The mixture was stirred at room temperature for 18 h and then a saturated solution of sodium bicarbonate (5 ml) was added. The product was extracted with ethyl acetate (3 x 2 ml) and the combined organic extracts were washed with brine, dried and concentrated *in vacuo* to leave a solid which was purified by preparative HPLC to give the sulfone, Compound 6 (10.0 mg, 46 %) as an off-white solid, δ<sub>H</sub> (400 MHz, MeOD) 8.11 (2H, obs dd *J* 7.9, 2.8 Hz, *Ar*), 7.79 (1H, dd *J* 9.6, 2.5 Hz, *Ar*), 7.65-7.57 (2H, m, *Ar*), 7.43 (1H, dd *J* 8.8, 4.3 Hz, *Ar*), 7.06 (1H, td *J* 9.1, 2.5 Hz, *Ar*), 4.76 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.85 (3H, s, CCH<sub>3</sub>); Tr =

25

30

1.44 min (100 %),  $m/z$  ( $ES^+$ ) ( $M+H$ ) $^+$  405.21, and the sulfoxide, Compound 7 (3.2 mg, 15 %) as an off-white solid,  $\delta_H$  (400 MHz, MeOD) 8.16 (1H, app d  $J$  9.1 Hz, *Ar*), 8.01 (1H, d  $J$  8.1 Hz, *Ar*), 7.62-7.54 (2H, m, *Ar*), 7.47 (1H, dd  $J$  9.1, 4.0 Hz, *Ar*), 7.23 (1H, dd  $J$  9.6, 2.5 Hz, *Ar*), 7.02 (1H, td  $J$  9.1, 2.0 Hz, *Ar*), 5.10 (2H, s,  $CH_2CO_2H$ ), 2.78 (3H, s,  $CCH_3$ ); Tr = 1.34 min (100 %),  $m/z$  ( $ES^+$ ) ( $M+H$ ) $^+$  389.09.

Compounds 8 to 10 were prepared using the same general method as for Compounds 6 and 7, but with appropriately chosen starting materials.

**10 Compound 8 – [5-Fluoro-2-methyl-3-(quinoline-2-sulfonyl)-indol-1-yl]-acetic acid**

$\delta_H$  (400 MHz, MeOD) 8.57 (1H, d  $J$  8.6 Hz, *Ar*), 8.20 (1H, d  $J$  8.6 Hz, *Ar*), 8.13 (1H, d  $J$  8.6 Hz, *Ar*), 8.02 (1H, d  $J$  8.1 Hz, *Ar*), 7.89-7.82 (2H, m, *Ar*), 7.73 (1H, app t  $J$  8.1 Hz, *Ar*), 7.42 (1H, dd  $J$  8.8, 4.3 Hz, *Ar*), 7.05 (1H, td  $J$  9.1, 2.5 Hz, *Ar*), 5.08 (2H, s,  $CH_2CO_2H$ ), 2.86 (3H, s,  $CCH_3$ ); Tr = 1.39 min (92 %),  $m/z$  ( $ES^+$ ) ( $M+H$ ) $^+$  399.26.

**Compound 9 – [5-Fluoro-2-methyl-3-(quinolin-8-ylsulfonyl)-indol-1-yl]-acetic acid**

$\delta_H$  (400 MHz, MeOD) 8.89 (1H, app d  $J$  4.3 Hz, *Ar*), 8.71 (1H, dd  $J$  7.3 Hz, *Ar*), 8.34 (1H, app d  $J$  8.3 Hz, *Ar*), 8.20 (1H, app d  $J$  8.3 Hz, *Ar*), 7.80 (1H, t  $J$  8.1 Hz, *Ar*), 7.58 (1H, dd  $J$  10.1, 2.5 Hz, *Ar*), 7.53 (1H, dd  $J$  8.3, 4.3 Hz, *Ar*), 7.34 (1H, dd  $J$  8.8, 4.3 Hz, *Ar*), 6.95 (1H, td  $J$  9.1, 2.5 Hz, *Ar*), 5.02 (2H, s,  $CH_2CO_2H$ ), 2.97 (3H, s,  $CCH_3$ ); Tr = 1.78 min (100 %),  $m/z$  ( $ES^+$ ) ( $M+H$ ) $^+$  399.29.

**25 Compound 10 – (5-Fluoro-2-methyl-3-phenylmethanesulfonyl-1H-indol-1-yl)-acetic acid**

$\delta_H$  (250 MHz,  $d_6$ -DMSO) 7.61 (1H, dd  $J$  9.0, 4.5 Hz, *Ar*), 7.35 (1H, dd  $J$  9.8, 2.5 Hz, *Ar*), 7.30-7.19 (3H, m, *Ar*), 7.10 (1H, td  $J$  9.1, 2.6 Hz, *Ar*), 7.02 (2H, m, *Ar*), 5.10 (2H, s,  $CH_2CO_2H$ ), 4.51 (2H, s,  $ArCH_2$ ), 2.06 (3H, s,  $CH_3$ ); Tr = 1.30min (100%)  $m/z$  ( $ES^+$ ) ( $M+H$ ) $^+$  362.13.

**Example 4 – Synthesis of 3-Sulfamoyl indole Derivatives (Method C)**

The method described below is employed for compounds of general formula (I) in which X is NR<sup>9</sup>.

5

**1. [3-(4-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid ethyl ester**

Chlorosulfonic acid (0.042 ml, 0.63 mmol) was added dropwise over 1 min to a stirred solution of (5-fluoro-2-methyl-indol-1-yl)-acetic acid ethyl ester (100 mg, 0.43 mmol) in ether (1 ml) at 0 °C. The solution was stirred at 0 °C for 10 min and then concentrated *in vacuo* to leave a residue which was azeotroped with dichloromethane (2 x 2 ml). The residue was taken up in dichloromethane and then *N,N*-diisopropyl ethylamine (0.075 ml, 0.43 mmol) and 4-chloroaniline (53.4 mg, 0.42 mmol) were added. The resulting mixture was stirred at room temperature for 40 min and then concentrated *in vacuo* to leave a residue which was partitioned between ethyl acetate (5 ml) and water (5 ml). The organic layer was then separated, washed with a saturated solution of sodium hydroxide (20 ml), dried and concentrated *in vacuo* to leave a residue which was purified by flash column chromatography (Flashmaster) on silica gel eluting with 15 % ethyl acetate : heptane to give the *sulfonamide* (6 mg, 3%) as an off-white solid,  $\delta_H$  (400 MHz, CDCl<sub>3</sub>) 7.63 (1H, dd *J* 9.5, 2.4 Hz, *Ar*), 7.18-7.12 (3H, m, *Ar*), 7.05-6.99 (1H, m, *Ar*), 6.96-6.90 (2H, m, *Ar*), 6.55 (1H, s, *NH*), 4.73 (2H, s, *NCH*<sub>2</sub>), 4.20 (2H, q *J* 7.3 Hz, *OCH*<sub>2</sub>*CH*<sub>3</sub>), 2.33 (3H, s, *CCH*<sub>3</sub>), 1.22 (3H, t *J* 7.3 Hz, *OCH*<sub>2</sub>*CH*<sub>3</sub>); *Tr* = 1.57 min (100 %), *m/z* (*ES*<sup>+</sup>) (*M*+*H*)<sup>+</sup> 425.

25

**2. Compound 11 – [3-(4-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

Lithium hydroxide monohydrate (7.0 mg, 0.17 mmol) in water (2 ml) was added in one portion to a stirred solution of [3-(4-chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid ethyl ester (6 mg, 0.014 mmol) in tetrahydrofuran (2 ml). The resulting mixture was stirred at room temperature for 3 h and then the pH of the

30

mixture was adjusted to pH 1 with 1M hydrochloric acid. The product was extracted with ethyl acetate (2 x 10 ml) and the combined organic extracts were then dried and concentrated *in vacuo* to give the *carboxylic acid* (4.3 mg, 77 %) as an off-white solid,  $\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 8.74 (1H, s, *NH*), 7.70 (1H, dd *J* 9.5, 2.6 Hz, *Ar*), 7.13-7.06 (3H, m, *Ar*), 6.99-6.92 (3H, m, *Ar*), 4.67 (2H, s,  $\text{NCH}_2$ ), 2.41 (3H, s,  $\text{CH}_3$ ); Tr = 1.84 min (91 %),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 397.

Compounds 12 to 25 were prepared using the same general method but with appropriately chosen starting materials.

10

**Compound 12 – [3-(3-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 7.63 (1H, dd *J* 9.3, 2.6 Hz, *Ar*), 7.17-7.14 (1H, m, *Ar*), 7.10-6.98 (5H, m, *Ar*, *NH*), 6.86-6.84 (1H, m, *Ar*), 4.73 (2H, s,  $\text{NCH}_2$ ), 2.46 (3H, s,  $\text{CH}_3$ ); Tr = 1.84 min (100 %),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 397.

15

**Compound 13 – [3-(4-Fluoro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 8.45 (1H, s, *NH*), 7.66 (1H, dd *J* 9.7, 2.3 Hz, *Ar*), 7.11 (1H, dd *J* 9, 4.2 Hz, *Ar*), 6.97-6.90 (3H, m, *Ar*), 6.81-6.77 (2H, m, *Ar*), 4.64 (2H, s,  $\text{NCH}_2$ ), 2.29 (3H, s,  $\text{CH}_3$ ); Tr = 1.79 min (99 %),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 381.

20

**Compound 14 – [3-(2-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $\text{CDCl}_3$ ) 7.69 (1H, s, *NH*), 7.58-7.49 (2H, m, *Ar*), 7.23-7.13 (3H, m, *Ar*), 7.03-6.93 (2H, m, *Ar*), 4.70 (2H, s,  $\text{NCH}_2$ ), 2.44 (3H, s,  $\text{CH}_3$ ); Tr = 1.83 (100 %),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ )<sup>+</sup> 397.

25

**Compound 15 – (3-Benzylsulfamoyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 7.99 (1H, t *J* 6.3 Hz, *Ar*), 7.58 (2H, m, *Ar*), 7.21 (4H, m, *Ar*), 7.09 (1H, td *J* 9.23, 2.65 Hz, *Ar*), 5.11 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 3.92 (2H, d *J* 6.31 Hz,

30

NCH<sub>2</sub>) 2.56 (3H, s, CH<sub>3</sub>), Tr = 1.31min (100%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 377.25.

**Compound 16 – [5-Fluoro-3-(2-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid**

5  $\delta_H$  (400 MHz, *d*<sub>6</sub>-DMSO) 9.18 (1H, s, SO<sub>2</sub>NH), 7.52 (1H, dd *J* 9.00, 4.4 Hz, *Ar*), 7.47 (1H, dd *J* 10.2, 2.6 Hz, *Ar*), 7.23 (1H, dd *J* 7.9, 1.6 Hz, *Ar*), 7.09-7.01 (2H, m, *Ar*), 6.84 (1H, t *J* 7.7 Hz, *Ar*), 6.77 (1H, d *J* 7.2 Hz, *Ar*), 5.05 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 3.24 (3H, OCH<sub>3</sub>), 2.30 (3H, s, CH<sub>3</sub>); Tr = 1.31 min (100%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup>  
10 393.25.

**Compound 17 – [5-Fluoro-3-(4-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid**

15  $\delta_H$  (400 MHz, *d*<sub>6</sub>-DMSO) 9.62 (1H, s, SO<sub>2</sub>NH), 7.49 (1H, dd *J* 10.3, 2.6 Hz, *Ar*), 7.36 (1H, dd *J* 9.0, 4.6 Hz, *Ar*), 6.98 (1H, dd *J* 9.2, 2.8 Hz, *Ar*), 6.93 (2H, d *J* 9.1 Hz, *Ar*), 6.74 (2H, d *J* 9.1 Hz, *Ar*), 4.45 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 3.64 (3H, s, OCH<sub>3</sub>), 2.36 (3H, s, CH<sub>3</sub>); Tr = 1.27 min (100%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 393.26.

**Compound 18 – (5-Fluoro-2-methyl-3-phenylsulfamoyl-indol-1-yl)-acetic acid**

20  $\delta_H$  (400 MHz, *d*<sub>6</sub>-DMSO) 10.15 (1H, s, SO<sub>2</sub>NH), 7.60 (1H, dd *J* 10.1, 2.6 Hz, *Ar*), 7.53 (1H, dd *J* 9.1, 4.5 Hz, *Ar*), 7.17 (2H, m, *Ar*), 7.07 (1H, dd *J* 9.1, 2.6 Hz, *Ar*), 7.03 (2H, m, *Ar*), 6.96 (1H, t *J* 7.3 Hz, *Ar*), 5.03 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.48 (3H, s, CH<sub>3</sub>); Tr = 1.28 min (96%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 363.25.

**Compound 19 – [3-(3,4-Dichloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

25  $\delta_H$  (400 MHz, *d*<sub>6</sub>-DMSO) 7.92 (1H, bs, SO<sub>2</sub>NH), 7.47 (1H, dd *J* 10.3, 2.6 Hz, *Ar*), 7.40 (1H, d *J* 8.3 Hz, *Ar*), 7.37 (1H, d *J* 1.7 Hz, *Ar*), 7.31 (1H, dd *J* 9.0, 4.6 Hz, *Ar*), 7.13 (1H, dd *J* 8.2, 2.0 Hz, *Ar*), 6.96 (1H, td *J* 9.3, 2.7 Hz, *Ar*), 4.32 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 3.93 (2H, s, NCH<sub>2</sub>), 2.49 (3H, s, CH<sub>3</sub>); Tr = 1.43 min (97%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 445.15.  
30

**Compound 20 – [5-Fluoro-3-(3-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 10.22 (1H, s,  $\text{SO}_2\text{NH}$ ), 7.60 (1H, dd  $J$  9.9, 2.5 Hz,  $\text{Ar}$ ), 7.50 (1H, dd  $J$  9.1, 4.1 Hz,  $\text{Ar}$ ), 7.05 (2H, m,  $\text{Ar}$ ), 6.61 (2H, m,  $\text{Ar}$ ), 6.50 (1H, d 8.7 Hz,  $\text{Ar}$ ), 4.88 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 3.60 (3H, s,  $\text{OCH}_3$ ), 2.54 (3H, s,  $\text{CH}_3$ ); Tr = 1.28 min (100%),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ ) $^+$  393.28.

**Compound 21 – (5-Fluoro-2-methyl-3-*m*-tolylsulfamoyl-indol-1-yl)-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 10.11 (1H, s,  $\text{SO}_2\text{NH}$ ), 7.61 (1H, dd  $J$  10.1, 2.6 Hz,  $\text{Ar}$ ), 7.55 (1H, dd  $J$  9.1, 4.5 Hz,  $\text{Ar}$ ), 7.08 (1H, dd  $J$  9.2, 2.6 Hz,  $\text{Ar}$ ), 7.03 (1H, d  $J$  7.8,  $\text{Ar}$ ), 6.85-6.81 (2H, m,  $\text{Ar}$ ), 6.76 (1H, d  $J$  7.6 Hz,  $\text{Ar}$ ), 5.08 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 2.50 (3H, s,  $\text{CH}_3$ ), 2.15 (3H, s,  $\text{ArCH}_3$ ), Tr = 1.33 min (100%),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ ) $^+$  377.24.

**Compound 22 – (5-Fluoro-2-methyl-3-*p*-tolylsulfamoyl-indol-1-yl)-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 9.90 (1H, bs,  $\text{SO}_2\text{NH}$ ), 7.55 (1H, dd  $J$  10.2, 2.6 Hz,  $\text{Ar}$ ), 7.32 (1H, dd  $J$  9.1, 4.6,  $\text{Ar}$ ), 6.94 (5H, m,  $\text{Ar}$ ), 4.31 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 2.45 (3H, s,  $\text{CH}_3$ ), 2.15 (3H, s,  $\text{CH}_3$ ), Tr = 1.33 min (100%),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ ) $^+$  377.25.

**Compound 23 – [3-(4-Chloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 8.04 (1H, t  $J$  5.9 Hz,  $\text{NH}$ ), 7.58-7.51 (2H, m,  $\text{Ar}$ ), 7.25 (2H, d  $J$  8.6 Hz,  $\text{Ar}$ ), 7.18 (2H, d  $J$  8.6 Hz,  $\text{Ar}$ ), 7.07 (1H, td  $J$  9.5, 2.7 Hz,  $\text{Ar}$ ), 5.07 (2H, s,  $\text{CH}_2\text{CO}_2\text{H}$ ), 3.93 (2H, d  $J$  6.3 Hz,  $\text{NCH}_2$ ), 2.56 (3H, s,  $\text{CH}_3$ ); Tr = 1.38 min (91%),  $m/z$  ( $\text{ES}^+$ ) ( $\text{M}+\text{H}$ ) $^+$  411.07.

**Compound 24 – [3-(Benzyl-methyl-sulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid**

$\delta_{\text{H}}$  (400 MHz,  $d_6$ -DMSO) 7.52 (1H, dd  $J$  10.0, 2.6 Hz,  $\text{Ar}$ ), 7.46 (1H, dd  $J$  8.9, 4.4



Hz, *Ar*), 7.39-7.28 (5H, m, *Ar*), 7.03 (1H, td *J* 9.4, 2.8 Hz, *Ar*), 4.46 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 4.10 (2H, s, NCH<sub>2</sub>), 2.60 (3H, s, NCH<sub>3</sub>), 2.48 (3H, s, CH<sub>3</sub>); Tr = 1.43 min (100%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 391.15.

5 **Compound 25 – [5-Fluoro-2-methyl-3-(pyridin-3-ylsulfamoyl)-indol-1-yl]-acetic acid**

$\delta_H$  (400 MHz, *d*<sub>6</sub>-DMSO) 10.42 (1H, s, SO<sub>2</sub>NH), 8.25 (1H, d *J* 2.6 Hz, *Ar*), 8.19 (1H, d, 3.3 Hz, *Ar*), 7.59-7.55 (2H, m, *Ar*), 7.40 (1H, d *J* 8.3 Hz, *Ar*), 7.24-7.21 (2H, m, *Ar*), 5.10 (2H, s, CH<sub>2</sub>CO<sub>2</sub>H), 2.49 (3H, s, CH<sub>3</sub>); Tr = 0.96 min (100%), *m/z* (ES<sup>+</sup>) (M+H)<sup>+</sup> 364.1.

10

**Example 5 – Measurement of CRTH2 Antagonist Activity**

**Materials and Methods**

15

**Materials**

Calcium-3 dye was purchased from Molecular Devices (Wokingham, UK). Monopoly resolving medium was obtained from Dainippon Pharmaceuticals (Osaka, Japan). Macs anti-CD16 microbeads were from Miltenyi biotec (Bisley, Surrey). ChemoTx plates were purchased from Neuroprobe (Gaithersburg, MD). Poly-D-lysine coated 96-well plates were obtained from Greiner (Gloucestershire, UK). [<sup>3</sup>H]PGD<sub>2</sub> was from Amersham Biosciences (Buckinghamshire, UK). [<sup>3</sup>H]SQ29548 was purchased from Perkin Elmer Life Sciences (Buckinghamshire, UK). All other reagents were obtained from Sigma-Aldrich (Dorset, UK), unless otherwise stated.

20

25

**Methods**

*Cell culture*

Chinese Hamster Ovary cells were transfected with CRTH2 or DP receptors (CHO/CRTH2 and CHO/DP) and were maintained in culture in a humidified atmosphere at 37°C (5% CO<sub>2</sub>) in Minimum Essential Medium (MEM) supplemented

30

with 10% foetal bovine serum, 2 mM glutamine, and 1 mg ml<sup>-1</sup> active G418. The cells were passaged every 2-3 days. For radioligand binding assay, cells were prepared in triple-layer flasks or in 175 cm<sup>2</sup> square flasks (for membrane preparation). For calcium mobilisation assay, cells were grown in a 96 well plate  
5 24h prior to the assay at a density of 80,000 cells per well.

#### *Preparation of cell membranes*

Membranes were prepared either from CHO/CRTH2 and CHO/DP cells, or from platelets (as a source of TP receptors). CHO cells grown to confluency were washed  
10 with PBS and detached using a Versene solution (15 ml per flask). When the cells were grown in 175 cm<sup>2</sup> square flask, they were collected by scrapping in PBS. The cell suspensions were centrifuged (1,700 rpm, 10 min, 4°C) and resuspended in 15 ml of buffer (1xHBSS, supplemented with 10 mM HEPES, pH 7.3). Cell suspensions were then homogenised using an Ultra Turrax at setting 4-6 for 20 s.  
15 The homogenate was centrifuged at 1,700 rpm for 10 min and the supernatant was collected and centrifuged at 20,000 rpm for 1h at 4°C. The resulting pellet was resuspended in buffer and stored at -80°C in aliquots of 200-500 µl. The protein concentration was determined by the method of Bradford (1976), using bovine serum albumin as standard. The platelets were washed by centrifugation at 600xg for 10  
20 min and resuspended in ice-cold assay buffer (10 mM Tris-HCl, pH 7.4, 5 mM Glucose, 120 mM NaCl, 10 µM indomethacin) and directly centrifuged at 20,000 rpm for 30 min at 4°C. The resulting pellet was treated as described above.

#### *Radioligand binding assays*

25 [<sup>3</sup>H]PGD<sub>2</sub> (160 Ci/mmol) binding experiments were performed on membranes prepared as described above. Assays were performed in a final volume of 100 µl of buffer (1XHBSS/HEPES 10 mM, pH 7.3). Cell membranes (15µg). Cell membranes 15mg were preincubated at room temperature with varying concentration of competing ligand for 15 min. [<sup>3</sup>H]PGD<sub>2</sub> (mol, final concentration) was then added  
30 and the incubation continued for a further one hour at room temperature. The reaction was terminated by the addition of 200 µl ice-cold assay buffer to each well,

followed by rapid filtration through Whatman GF/B glass fibre filters using a Unifilter Cell harvester (PerkinElmer Life Sciences) and six washes of 300  $\mu$ l of ice-cold buffer. The Unifilter plates were dried at room temperature for at least 1h and the radioactivity retained on the filters was determined on a Beta Trilux counter  
5 (PerkinElmer Life Sciences), following addition of 40  $\mu$ l of Optiphase Hi-Safe 3 (Wallac) liquid scintillation. Non specific binding was defined in the presence of 10  $\mu$ M unlabelled PGD<sub>2</sub>. Assays were performed in duplicate.

The results of the radioligand binding experiments to the CRTH2 and DP receptors  
10 are shown in Table 1.

The results shown in Table 1 demonstrate that for compounds of general formula (I) have high affinity for the CRTH2 receptor. In those cases where a comparison was made, the affinity of the compounds of general formula (I) is much higher for the  
15 CRTH2 receptor than for DP receptor.

**Table 1 – Radioligand binding data (K<sub>i</sub> on CRTH2 Receptor and DP Receptor).**

<b><u>Compound</u></b>	<b>CRTH2 Binding K<sub>i</sub> nM</b>	<b>DP Binding K<sub>i</sub> <math>\mu</math>M</b>
1	192	>10
2	75	>10
3	2000	ND
4	2300	ND
5	89	>10
6	209	ND
7	54	ND
8	249	ND
9	254	>10
10	6	ND
11	51	>10
12	45	>10
13	182	ND
14	225	ND
15	278	>10
16	771	>10
17	1450	>10
18	236	>10
19	181	>10
20	531	>10
21	176	>10
22	1240	>10
23	164	>10
24	119	>10
25	4250	>10

- 5 The TP receptor radioligand binding was done on membranes prepared from platelets. 15-40  $\mu$ g of protein were pre-incubated with varying concentrations of competing ligand for 15 min at room temperature in assay buffer (10 mM Tris-HCl, pH 7.4, 5 mM glucose, 120 mM NaCl, 10  $\mu$ M indomethacin). [<sup>3</sup>H]SQ29548 (38 Ci/mmol, 10 nM final concentration) was then added and the incubation continued
- 10 for a further 30 min at room temperature. The reaction was terminated by the addition of 200  $\mu$ l ice-cold assay buffer to each well, followed by rapid filtration through Whatman GF/C glass fibre filters using a Unifilter Cell harvester (PerkinElmer Life Sciences) followed with six washes of 300  $\mu$ l of ice-cold buffer.

The radioactivity was determined as described above.

All of the compounds studied in this assay bound to the TP receptor with low affinity ( $K_i > 10 \mu\text{M}$ ).

5

Compounds of general formula (I) bound to CRTH2 receptor expressed in CHO cells with a range of affinity varying from very high to moderate. In fact the  $K_i$  values determined in competition versus [ $^3\text{H}$ ]PGD<sub>2</sub> varied from 500 pM to 1  $\mu\text{M}$ . Compounds of general formula (I) had no activity (or very weak activity) at the DP and TP receptors. The binding selectivity of the compounds of general formula (I) for CRTH2 receptor was greater than 200 fold for CRTH2 receptor, compared to DP and TP receptors.

10

#### *Calcium mobilisation Assay*

15

Cells were seeded onto poly-D-lysine coated 96-well plates at a density of 80,000 cells per well and incubated at 37°C overnight to allow the cells to adhere. Cells were washed twice with HBSS and incubated for 1h at 37°C in 100 $\mu\text{l}$  HBSS and 100 $\mu\text{l}$  calcium-3-dye (Molecular Devices) solution, supplemented with 4mM probenecid. Changes in fluorescence were monitored over a 50s time course with agonist addition at 17s using a Flexstation (Molecular Devices).

20

#### *Effect of CRTH2 agonists on calcium mobilisation in CHO-CRTH2 cells*

PGD<sub>2</sub> caused a dose-dependent increase in intracellular  $\text{Ca}^{2+}$  mobilisation in CHO/CRTH2 cells, with an  $\text{EC}_{50} = 2.4 \pm 0.5 \text{ nM}$  ( $n=3$ ) (Figure 1).

25

#### *Effect of compounds of general formula (I) on the calcium mobilisation induced by PGD<sub>2</sub>*

PGD<sub>2</sub>-stimulated  $\text{Ca}^{2+}$  flux was fully inhibited by the compounds of general formula (I) and the  $\text{IC}_{50}$  value for each compound in the calcium assay was comparable to its  $K_i$  value in Radioligand binding.  $\text{IC}_{50}$  values of compounds of general formula (I) varied from 5 nM to 1  $\mu\text{M}$ . The results for several compounds of general formula (I) are shown in Table 2. Increasing doses of the compounds of general formula (I) caused a dose-dependent and parallel shift of the PGD<sub>2</sub> dose response curve in

30

CHO/CRTH2 cells, thereby indicating that the compounds are competitive CRTH2 antagonists.


The antagonistic effect of the compounds of general formula (I) appears to be  
5 CRTH2 selective, since no inhibitory effect was seen with ATP-stimulated  $\text{Ca}^{2+}$  flux in CHO/CRTH2 cells.

**Table 2 – Inhibition of  $\text{PGD}_2$ -induced calcium flux**

Compound	CRTH2 Ca flux $\text{IC}_{50}$ (nM)
1	280
2	163
5	268
6	345
7	163
8	330
10	79
11	197
12	82
13	1650
14	390
21	1060

CLAIMS

1. A compound of general formula (I)



I

wherein

$R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are independently hydrogen, halo,  $C_1$ - $C_6$  alkyl,  $-O(C_1$ - $C_6$  alkyl),  $-CON(R^9)_2$ ,  $-SOR^9$ ,  $-SO_2R^9$ ,  $-SO_2N(R^9)_2$ ,  $-N(R^9)_2$ ,  $-NR^9COR^9$ ,  $-CO_2R^9$ ,  $-COR^9$ ,  $-SR^9$ ,  $-OH$ ,  $-NO_2$  or  $-CN$ ;

each  $R^9$  is independently hydrogen or  $C_1$ - $C_6$  alkyl;

$R^5$  and  $R^6$  are each independently hydrogen, or  $C_1$ - $C_6$  alkyl or together with the carbon atom to which they are attached form a  $C_3$ - $C_7$  cycloalkyl group;

$R^7$  is hydrogen or  $C_1$ - $C_6$  alkyl

$n$  is 1 or 2;

$X$  is a bond or, when  $n$  is 2,  $X$  may also be a  $NR^9$  group;

wherein  $R^9$  is as defined above;

when  $X$  is a bond  $R^8$  is  $C_1$ - $C_6$  alkyl,  $C_2$ - $C_6$  alkenyl,  $C_2$ - $C_6$  alkynyl, biphenyl or a 9-14 membered bicyclic or tricyclic heteroaryl group;

when  $X$  is a  $NR^9$  group  $R^8$  may additionally be phenyl, naphthyl or a 5-7 membered heteroaromatic ring; and

the  $R^8$  group is optionally substituted with one or more substituents selected from halo,  $C_1$ - $C_6$  alkyl,  $-O(C_1$ - $C_6$ )alkyl, aryl,  $-O$ -aryl, heteroaryl,  $-O$ -heteroaryl,  $-CON(R^9)_2$ ,  $-SOR^9$ ,  $-SO_2R^9$ ,  $SO_2N(R^9)_2$ ,  $-N(R^9)_2$ ,  $-NR^9COR^9$ ,  $-CO_2R^9$ ,  $-COR^9$ ,  $-SR^9$ ,


-OH, -NO<sub>2</sub> or -CN;

wherein R<sup>9</sup> is as defined above;

or a pharmaceutically acceptable salt, hydrate, solvate, complex or prodrug thereof.

5

2. A compound of general formula (II):



II

10 wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, n, X, R<sup>7</sup> and R<sup>8</sup> are as defined for general formula (I);  
R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, (CH<sub>2</sub>)<sub>m</sub>OC(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, (CH<sub>2</sub>)<sub>m</sub>N(R<sup>11</sup>)<sub>2</sub>,  
CH((CH<sub>2</sub>)<sub>m</sub>O(C=O)R<sup>12</sup>)<sub>2</sub>;

m is 1 or 2;

R<sup>11</sup> is hydrogen or methyl;

15 R<sup>12</sup> is C<sub>1</sub>-C<sub>18</sub> alkyl.

3. A compound as claimed in claim 1 or claim 2 wherein, independently or in any combination:

R<sup>1</sup> is halo or hydrogen;

20 R<sup>2</sup> is halo or hydrogen;

R<sup>3</sup> is halo or hydrogen;

R<sup>4</sup> is halo or hydrogen.

4. A compound as claimed in any one of claims 1 to 3 wherein R<sup>1</sup>, R<sup>3</sup> and R<sup>4</sup> are  
25 hydrogen and R<sup>2</sup> is halo.



5. A compound as claimed in claim 4 wherein  $R^2$  is fluoro.
6. A compound as claimed in any one of claims 1 to 5 wherein  $R^5$  and  $R^6$  are  
5 each independently hydrogen or  $C_1$ - $C_4$  alkyl.
7. A compound as claimed in claim 6 wherein at least one of  $R^5$  and  $R^6$  are hydrogen.
- 10 8. A compound as claimed in claim 7 wherein both  $R^5$  and  $R^6$  are hydrogen.
9. A compound as claimed in any one of claims 1 to 8 wherein  $R^7$  is H or  $C_1$ - $C_6$  alkyl.
- 15 10. A compound as claimed in claim 9 wherein  $R^7$  is methyl.
11. A compound as claimed in any one of claims 1 to 10 wherein n is 2.
12. A compound as claimed in any one of claims 1 to 11 wherein X is a bond and  
20  $R^8$  is  $C_1$ - $C_6$  alkyl, biphenyl or a bicyclic heteroaryl group, any of which may be substituted with halogen, phenyl,  $-CO_2R^9$   $CON(R^9)_2$  or  $-SO_2R^9$ , where  $R^9$  is as defined above.
13. A compound as claimed in claim 12 wherein  $R^8$  is  $C_1$ - $C_4$  alkyl, biphenyl, a  
25 bicyclic heteroaryl group or a 5-7 membered heterocyclic ring, any of which may be substituted with phenyl,  $-CO_2R^9$   $CON(R^9)_2$  or  $-SO_2R^9$ , where  $R^9$  is H or  $C_1$ - $C_4$  alkyl.
14. A compound as claimed in any one of claims 1 to 11 wherein X is  $NR^9$ ,  $R^9$  is H or methyl and  $R^8$  is:  
30 phenyl optionally substituted with one or more halo,  $C_1$ - $C_6$  alkyl or  $-O(C_1$ - $C_6$  alkyl) groups;

C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted with aryl; or  
heteroaryl.

15. A compound as claimed in claim 14, wherein R<sup>8</sup> is phenyl, benzyl or pyridyl,  
5 any of which may optionally be substituted with one or more halo, methyl or methoxy groups.
16. [3-(Butane-1-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
3-(Biphenyl-4-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 10 (3-Carboxymethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid  
(3-Carbamoylmethanesulfonyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid  
[5-Fluoro-3-(2-methanesulfonyl-ethanesulfonyl)-2-methyl-indol-1-yl]-acetic acid  
[3-(Benzothiazole-2-sulfonyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
[3-(Benzothiazole-2-sulfinyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 15 [5-Fluoro-2-methyl-3-(quinoline-2-sulfonyl)-indol-1-yl]-acetic acid  
[5-Fluoro-2-methyl-3-(quinolin-8-ylsulfonyl)-indol-1-yl]-acetic acid  
(5-Fluoro-2-methyl-3-phenylmethanesulfonyl-1H-indol-1-yl)-acetic acid  
[3-(4-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
[3-(3-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid
- 20 [3-(4-Fluoro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
[3-(2-Chloro-phenylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
(3-Benzylsulfamoyl-5-fluoro-2-methyl-indol-1-yl)-acetic acid  
[5-Fluoro-3-(2-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid  
[5-Fluoro-3-(4-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid
- 25 (5-Fluoro-2-methyl-3-phenylsulfamoyl-indol-1-yl)-acetic acid  
[3-(3,4-Dichloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
[5-Fluoro-3-(3-methoxy-phenylsulfamoyl)-2-methyl-indol-1-yl]-acetic acid  
(5-Fluoro-2-methyl-3-*m*-tolylsulfamoyl-indol-1-yl)-acetic acid  
(5-Fluoro-2-methyl-3-*p*-tolylsulfamoyl-indol-1-yl)-acetic acid
- 30 [3-(4-Chloro-benzylsulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid  
[3-(Benzyl-methyl-sulfamoyl)-5-fluoro-2-methyl-indol-1-yl]-acetic acid

[5-Fluoro-2-methyl-3-(pyridin-3-ylsulfamoyl)-indol-1-yl]-acetic acid;  
 or the C<sub>1</sub>-C<sub>6</sub> alkyl, aryl, (CH<sub>2</sub>)<sub>m</sub>OC(=O)C<sub>1</sub>-C<sub>6</sub>alkyl, (CH<sub>2</sub>)<sub>m</sub>N(R<sup>11</sup>)<sub>2</sub>,  
 CH((CH<sub>2</sub>)<sub>m</sub>O(C=O)R<sup>12</sup>)<sub>2</sub> esters of any of the above; wherein

m is 1 or 2;

5 R<sup>11</sup> is hydrogen or methyl;

R<sup>12</sup> is C<sub>1</sub>-C<sub>18</sub> alkyl.

17. A process for the preparation of a compound of general formula (I) as  
 claimed in any one of claims 1 to 13 or 16 wherein n is 1 or 2 and X is a bond, the  
 10 process comprising treating a compound of general formula (Ia), which is a  
 compound of general formula (I) wherein n is 0 and X is a bond, by oxidation with a  
 suitable oxidising agent.

18. A process for the preparation of a compound of general formula (I) as  
 15 claimed in any one of claims 1 to 16, the process comprising reacting a compound of  
 general formula (II) as defined in claim 2 and wherein R<sup>10</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl with a base.

19. A compound as claimed in any one of claims 1 to 16 for use in medicine.

20. A compound for use in the treatment of allergic asthma, perennial allergic  
 rhinitis, seasonal allergic rhinitis, atopic dermatitis, contact hypersensitivity  
 (including contact dermatitis), conjunctivitis, especially allergic conjunctivitis,  
 eosinophilic bronchitis, food allergies, eosinophilic gastroenteritis, inflammatory  
 bowel disease, ulcerative colitis and Crohn's disease, mastocytosis, another PGD<sub>2</sub>-  
 25 mediated disease, for example autoimmune diseases such as hyper IgE syndrome and  
 systemic lupus erythematus, psoriasis, acne, multiple sclerosis, allograft rejection,  
 reperfusion injury and chronic obstructive pulmonary disease; or rheumatoid  
 arthritis, psoriatic arthritis or osteoarthritis.

30 21. The use of a compound as claimed in any one of claims 1 to 16 in the  
 preparation of an agent for the treatment or prevention allergic asthma, perennial

- allergic rhinitis, seasonal allergic rhinitis, atopic dermatitis, contact hypersensitivity (including contact dermatitis), conjunctivitis, especially allergic conjunctivitis, eosinophilic bronchitis, food allergies, eosinophilic gastroenteritis, inflammatory bowel disease, ulcerative colitis and Crohn's disease, mastocytosis, another PGD<sub>2</sub>-mediated disease, for example autoimmune diseases such as hyper IgE syndrome and systemic lupus erythematus, psoriasis, acne, multiple sclerosis, allograft rejection, reperfusion injury and chronic obstructive pulmonary disease; or rheumatoid arthritis, psoriatic arthritis or osteoarthritis.
- 10 22. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 16 together with a pharmaceutical excipient or carrier.
- 15 23. A composition as claimed in claim 22 formulated oral, rectal, nasal, bronchial (inhaled), topical (including eye drops, buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration.
24. A composition as claimed in claim 23 formulated for oral, nasal, bronchial or topical administration.
- 20 25. A composition as claimed in any one of claims 22 to 24 containing one or more additional active agents useful in the treatment of diseases and conditions mediated by PGD<sub>2</sub> at the CRTH2 receptor.
- 25 26. A composition as claimed in claim 25, wherein the additional active agents are selected from:
- β<sub>2</sub> agonists such as salmeterol;
  - corticosteroids such as fluticasone;
  - antihistamines such as loratidine;
  - leukotriene antagonists such as montelukast;
  - 30 anti-IgE antibody therapies such as omalizumab;
  - anti-infectives such as fusidic acid (particularly for the treatment of atopic

dermatitis);

anti-fungals such as clotrimazole (particularly for the treatment of atopic dermatitis);  
immunosuppressants such as tacrolimus and particularly pimecrolimus in the case of  
inflammatory skin disease;

- 5 other antagonists of  $\text{PGD}_2$  acting at other receptors such as DP antagonists;  
inhibitors of phosphodiesterase type 4 such as cilionilast;  
drugs that modulate cytokine production such as inhibitors of  $\text{TNF}\alpha$  converting  
enzyme (TACE);  
drugs that modulate the activity of Th2 cytokines IL-4 and IL-5 such as blocking  
10 monoclonal antibodies and soluble receptors;  
PPAR- $\gamma$  agonists such as rosiglitazone;  
5-lipoxygenase inhibitors such as zileuton.


27. A process for the preparation of a pharmaceutical composition as claimed in  
15 any one of claims 22 to 26 comprising bringing a compound as claimed in any one of  
claims 1 to 16 in conjunction or association with a pharmaceutically or veterinarily  
acceptable carrier or vehicle.

28. A product comprising a compound as claimed in any one of claims 1 to 16  
20 and one or more of the agents listed in claim 26 as a combined preparation for  
simultaneous, separate or sequential use in the treatment of a disease or condition  
mediated by the action of  $\text{PGD}_2$  at the CRTH2 receptor.

29. The use as claimed in claim 21, wherein the agent also comprises an  
25 additional active agent useful for the treatment of diseases and conditions mediated  
by  $\text{PGD}_2$  at the CRTH2 and/or DP receptor.

30. The use as claimed in claim 29, wherein the additional active agent is one of  
the agents listed in claim 26.

1/1

**FIGURE 1**

15

# INTERNATIONAL SEARCH REPORT

PCT/GB2004/004336

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D209/30 A61K31/404 A61P43/00 C07D403/12 C07D401/12  
C07D417/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	IMMANUEL F. LÜSCHER ET AL.: "Deblocking of o-Nitrophenylsulfenyl-protected peptides by ammonium thiocyanate and (2-methyl-1-indolyl)acetic acid" HELVETICA CHIMICA ACTA., vol. 66, no. 2, 1983, pages 602-605, XP002269354 VERLAG HELVETICA CHIMICA ACTA. BASEL., CH ISSN: 0018-019X * page 604, compound 8 * ----- -/--	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*8\* document member of the same patent family

Date of the actual completion of the international search

12 January 2005

Date of mailing of the international search report

31/01/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Diederer, J

# INTERNATIONAL SEARCH REPORT

PCT/GB2004/004336

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOSEFINA GARCIA ET AL.: "A novel synthesis of 3-cyanoindoles and a new route to indole-3-carboxylic acid derivatives" TETRAHEDRON LETTERS., vol. 26, no. 15, 1985, pages 1827-1830, XP002269353 ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM., NL ISSN: 0040-4039 * page 1828, scheme I, compound 3 ( RN 98508-71-5 ) *	2
A	----- WO 03/066047 A (ASTRAZENECA AB) 14 August 2003 (2003-08-14) cited in the application page 1	1,21,22
P,X	----- WO 03/101961 A (ASTRAZENECA AB) 11 December 2003 (2003-12-11) * complete document *	1-11, 20-25,27
	-----	



# INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB2004/004336

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 03066047	A	14-08-2003	EP	1474137 A1	10-11-2004
			WO	03066047 A1	14-08-2003
WO 03101961	A	11-12-2003	WO	03101961 A1	11-12-2003